

ZARETSKII, B. I.

Tractors. Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit.lit-ry,
1950. 135 p. (50-55251)

TL233.Z3

KITAYEV, Ye.N., inzh.; ZARETSKIY, B.I., otv. red.

[The best cements for the manufacture of asbestos-cement materials] Optimal'nye tsementy dlia proizvodstva asbestotsementnykh materialov. Moskva, Otdel nauchno-tekhn.informatsii, 1959. 71 p.

(Cement) (Asbestos cement)

KITAYEV, Ye.N., inzh.; GONCHARSKAYA, R.E.; ZARETSKIY, B.I., ctv. red.;
ERLIKH, I.A., red.

[Asbestos cement materials obtained from sand cements by
autoclave treatment, and their chemical resistance to cor-
rosive solutions] Khimicheskaiia stoikost' v agressivnykh
rastvorakh asbestotsementnykh materialov, poluchaemykh iz
peschanistykh tsementov s primeneniem avtoklavnoi obrabotki.
Moskva, Otdel nauchno-tekhn. informatsii, 1960. 24 p.

(MIRA 15:1)

(Asbestos cement)

ZARETSKIY, B.I.

KAREL'SKIKH, D.K., prof.; APASHEV, M.D., kand.tekhn.nauk; BARSKIY, I.B.,
kand.tekhn.nauk; ZAYCHIK, G.I., doktor tekhn.nauk, retsenzent;
ANOKHIN, V.I., kand.tekhn.nauk, retsenzent; ZARETSKIY, B.I.,
inzh., red.; POPOVA, S.M., tekhn.red.

[Theory, design, and engineering analysis of tractors] Teoriia,
konstruktsiia i raachet traktorov. Moskva, Gos.nauchno-tekhn.
izd-vo mashinostroit. lit-ry. [Pt.3]. Theory and analysis of
tractor chassis] Teoriia i raachet shassi traktorov. Pod obshchei
red. D.K.Karel'skikh. 1950. 144 p. (MIRA 11:12)
(Tractors)

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1

ZARETSKIY, B. I.

"Tractors," Moscow 1950, 1 copy.

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1"

L 29931-66 EWP(k)/EWT(d)/EWT(m)/EWP(h)/T/EWP(1)/EWP(v)/EWP(t)/ETI JD/HM
ACC NR: AP6018011 (A) SOURCE CODE: UR/0413/66/000/010/0126/0126

INVENTOR: Voronin, G. I.; Slotin, V. I.; Zaretskiy, B. S.; Krylov, A. I.;
Shvetsov, P. N.; Barannikov, G. I.; Eskin, G. I.

4D

B

ORG: none

TITLE: Ultrasonic unit for fluxless brazing of metals. Class 49, No. 181967

SOURCE: Izobreteniya, promyshlennyye obraztsy, tovarnyye znaki, no. 10, 1966, 126

TOPIC TAGS: brazing, metal brazing, ultrasonic brazing, brazing unit

ABSTRACT: This Author Certificate introduces a unit for fluxless brazing of metals equipped with a heater and ultrasonic emitter. To increase efficiency, the ultrasonic




Fig. 1. Fluxless brazing unit

1 - Crucible; 2 - brazing alloy;
3 - ultrasonic emitter.

Card 1/2

UDC: 621.791.351.6.03

L 29931-66

ACC Nbr AP6018011

emitter is located inside the crucible containing molten brazing alloy, forming
the bottom of the latter (see Fig. 1.). Orig. art. has: 1, figure. [AZ]

SUB CODE: 11,13/SUBM DATE: 29Jan65/ ATD PRESS: 5011

Card 2/2 CC

BAT*, G.A.; ZARETSKIY, D.F.

[Solution of the generalized Milne problem] Reshenie obobshchennoi zadachi Milna. Minsk, 1955. 15 p. (MIR. 14:7)
(Neutrons—Scattering) (Integral equations)

ZARETSKIY, D.P.

[Effective boundary conditions for "gray" bodies] Efektivnye
granichnye usloviia dlia "serykh" tel; doklady, predstavленные
SSSR na Mezhdunarodnii konferentsii po mirnomu ispol'zovaniyu
atomnoi energii. Moskva, 1955. 17 p. [Microfilm]
(Nuclear physics) (MIRA 9:3)

ZARETSKIY
USSR/Nuclear Physics - Thermal neutron capture

FD-3260

Card 1/1 Pub. 146 - 19/44

Author : Shut'ko, A. V.; Zaretskiy, D. F.

Title : Capture of thermal neutrons by lead isotopes

Periodical : Zhur. eksp. i teor. fiz., 29, No 6(12), Dec 1955, 867-868

Abstract : The authors consider the isotopes Pb-207 and Pb-208 and their excitation levels, spins, parities, energies, etc. They compare the theoretical evaluations of cross-section of thermal neutron capture by lead isotopes with data of experiments. They conclude that capture in Pb-206 is "less single-particle" than in the case of Pb-207, and that the anomalous character of capture radiation in lead isotopes is explained by proceeding from the single-particle picture of capture. The authors thank Professor A. S. Davydov and V. F. Turchin for comments. Seven references, including one USSR: L. K. Peker and L. A. Sliv, Izv. AN SSSR, ser. fiz., 17, 1953.

Institution : --

Submitted : August 31, 1955

ZARETSKIY, D. F. and SHUTKO, A. V.

"On the Thermal Capture by Pb Isotopes" a paper presented at the International Conference on Nuclear Reactions, Amsterdam, 2-7 July 1956.

D551274

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1

Absorption of thermal neutrons by isotopes of lead
V. Saut'ev and D. F. Zaitsev. Sov. Phys., JETP, 2,
702-71(1956) (English translation). See C.A. 50, 14373.
B.M.R. Sci

2


Ref

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1"

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1



APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1"

AUTHOR ZARETSKIY, D.F., SHUTKO, A.V. PA - 2690
TITLE On the Quasi-Magnetic Interaction of the Nucleon Spin With the
Rotation of the Nucleus.
(O kvazimagniton vzaimodeystvii spina nuklona s vrashcheniyem yadra-
Russian)
PERIODICAL Zhurnal Eksperim. i Teoret. Fiziki, 1957, Vol 32, Nr 2, pp 370-371,
(U.S.S.R.)
Received 5/1957 Reviewed 6/1957
ABSTRACT It is possible to find a new interpretation for the rotational levels of
nuclei with spin 1/2 if we start out from the following premises: (1) In
These nuclei there exist Σ -states. Then in first approximation the levels
with the total angular momentum $I=K \pm 1/2$ are degenerated. (Here K stands
for the rotational quantum number). (2) This degeneration is eliminated
if we introduce into the Hamiltonian by Bohr and Mottelson an interaction
of the form of $H_{Rs} = -(\lambda/mc^2) \vec{S} \cdot [U(\vec{r}) \vec{v}_{koll}]$. Here λ stands for a non-
dimensional phenomenological constant with the same significance and ma-
gnitude as in the normal (usual) spin-orbit coupling of the nucleus.
Furthermore the following denotations are used: \vec{S} for the vector of the
nucleon spin. $U(\vec{r})$ for the selfconsisting potential of the nucleus, mc^2
for the rest energy of the nucleon. and v_{koll} for the velocity with which
the nucleon participates in the collective motion. First of all the sig-
nificance of v_{koll} is clarified, and then the significance and the ori-
gin of the above-mentioned interaction H_{Rs} . For the wave function we set
Card 1/2

AUTHORS:

Bart', G.A., Zarotskiy, D.F.

SOV/89-4-6-2/30

TITLE:

The Effective Boundary Value Conditions in the Diffusion Theory
of Neutrons (Survey) (Effektivnyye granichnyye usloviya v teorii
diffuzii neytronov (Obzor))

PERIODICAL:

Atomnaya energiya, 1958, Vol. 4, Nr 6, pp. 510-519 (USSR)

ABSTRACT:

In the course of a survey the methods are described by means of which the effective boundary value conditions can be determined which bring about agreement within the asymptotic range between the solution of the neutron diffusion equation and the solution of the corresponding kinetic equation.

The boundary values are described for monoenergetic neutrons; some of them are computed, and, especially, the occurring coefficients are registered in tables for the following cases:

- 1.) For an infinitely thin "black" rod in a medium, in which sources are uniformly distributed.
- 2.) For a plane boundary between medium and vacuum by means of the exact analytical method developed by Wiener and Hopf.
- 3.) For a "black" round cylinder. The following methods of solving are mentioned for this purpose:

Card 1/2

The Effective Boundary Value Conditions in the
Diffusion Theory of Neutrons

SOV/39-4-6-2/30

- a) The balance method developed by Brudno (1951)
- b) The variation method developed by Zaretskiy (Ref 6)
- c) The method of spherical harmonics by Galanin
- d) Solution of Peierls (Payerl's) equations for great and small r_0 according to the method developed by Davison.
- 4.) For a "black" rod of any cross section by the method of approximation developed by Hurwitz (Gurvits) and Roe (Ref 8).
- 5.) For a "gray" rod (only possible by approximation). If it is intended to take the neutron spectrum into account in connection with the boundary value conditions, this is possible only for some simple border cases.
(A body is described as being either black or gray according to whether the neutrons impinging upon its surface are fully or only partly absorbed). There are 2 figures, 5 tables, and 10 references, 4 of which are Soviet.

SUBMITTED:

October 4, 1957

1. Neutrons--Diffusion 2. Diffusion--Theory 3. Mathematics
--Applications

Card 2/2

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1

ZARETSKIY, D. F.

"On -Masonic Fission."

paper to be presented at 2nd Un Intl. Conf. on the peaceful uses of Atomic Energy, Geneva, 1 - 13 Sep 58.

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1"

DROZDOV, S. I., ZARETSKIY, D. F., KURRIN, L. P. and SEDELNIKOV, T. Kh.

"On the Formation of a Thermal Neutron Spectrum."

Paper to be presented at 2nd UN Ints. Conf. on the peaceful uses of Atomic Energy, Geneva, 1 - 13 Sep 58.

ZARETSKII, D.F.

INOPIN, Ye.V.

AUTHOR: Verhalovich, D.

SERIAL #: SCI/53-65-4-7/15

TITLE: The VIII Annual Congress of Nuclear Spectroscopy (VIII
Zashchegodzhe sverkhshchisly po Yadernoj spetskopeji). 2

PERIODICAL: Uspekhi fizicheskikh nauk, 1959, Vol. 65, No. 4.

pp. 721 - 722 (USSR)

ABSTRACT: The 8th Congress of Nuclear Spectroscopy took place in Leningrad from January 27 to February 3, 1959. It was attended by 300 scientists from the USSR, further by scientists from China, France, Poland, Czechoslovakia, Hungary, Germany, Yugoslavia, and the Federal Democratic Republic of民主德意志 and about 100 guests were invited. The main lectures dealt with problems concerning nuclear models, the α - and β -decay, γ -radiation, internal conversion, and nuclear fission. B.D. Dubleiko, Gor'kopol'skiy, Slobodanov, and many others opened the conference. Lectures were held by V.Ye. Gonorov, Ye.V. Inopin, G.I. Tsvetkov (TGU, AS USSR) on light nuclides and general nuclear models; Peter (Budapest University, AS USSR); Yu.M. Shirokov (BSC-Moscow State University), L.A. Leshin (LPI-Leningrad Physical-Techical Institute) on levels in ^{16}O , ^{18}O and ^{12}C ; D.G.

Aleksandrov, A.P. Zhitomirsky, G.M. Galatsky, E.I. Terolianskaia, T.N. Vinogradova (LPI) on having found no rotational levels at 2^+ in ^{40}Ca , ^{48}Ca , and ^{56}Ni . The same research workers also reported on the discovery of vibrational γ -levels in ^{16}O (2^+ , 3^+ , 4^+) nuclei by means of the method of the Coulomb (Kulon) excitation at $E_{\gamma} = 1$ MeV. L.K. Peker (BAS 252R) gave a survey report: "Concerning Some Features in Vibrational Levels of Deformed Nuclei." Lectures were held also by D.V. Zaretskii (IAE USSR - AIPER) on transitions in ^{16}O and ^{18}O nuclei on irradiation. Shibanov, T. MIFP (MIFI (Moscow State University) on the level displacements and the probability corresponding to β^- and γ^- transitions in odd nuclei; D. Zaretskii (IAE USSR - AS USSR) on the influence of the spin-orbital coupling upon the magnetic moments of the nucleus; A.F. Barin (IAE USSR - IAE USSR) on the existence of light nuclei with high neutron or proton excesses; V.I. Kravtsov (IUP-Berlin) on the formation of nucleon pairs in nuclei; L. A. Gordian, A. N. Pilyayev, G.M. Martiroyan, K.A.Tser-Martiroyan (IAE USSR).

on alpha decay on rotational levels of odd nuclei; V.G. Savor (IAE USSR - AS USSR) on alpha decay of nonpherical nuclei (survey); A.I. Alakhbarov, G.P. Veliev, V. V. Zhuravlev, V.Y. Zhuravlev (TGU, AS USSR) on measurements of electrons emitted in the β -decay of ^{12}C , ^{14}N , ^{15}O , ^{16}O ($0^+, 1^-, 2^-, 3^-$) as well as in that of ^{36}S and ^{38}Se ; F.A. Baryev and Yu. Zaretskii (TGU, AS USSR) on investigations of electron-electron correlations and the resonance scattering of e^-e^+ in ^{16}O ; A.Z. Karabut and V.M. Prishchepov (MIFI-Berlin State University) on the measurement of longitudinally polarized electrons; A.I. Zubarev and Yu. V. Perov (MCU) on the effect of the orientation of the cathode of polarized electrons and positions of the cathodes; Ya.E. Chudarev and I.Y. Faure (BIP) on polarized electrons and the intensity of the components of the complex $\beta-\gamma\gamma\gamma$ process according to the Fermi diagram; A.Z. Shao, L.I. Zytseva, andputation of the probability of the permitted and of the forbidden capture of electrons by a nucleus.

ZARETSKIY, D. F.

2. (b) **REPORTS**
 International Conference on the Chemical Basis of Acute Mortality, 2d, Odessa, 1958
 Soviet approach towards Industrial Safety (Report of Soviet Scientific Committee on Chemical Safety). Moscow, Academy of Sciences, 1970. 55 p. (Series: The Study, vol. 2; No. 20; English translation)

2. (b) **REPORTS**
 V.A. Savchenko, Candidate of Physic and Mathematical Sciences, Dr. of this
 Committee, B.I. Smirnov and B.P. Savchenko, Candidates of Physical and Mathematical
 Sciences Dr. (Candidate Doctoral Dissertation). Moscow, 1971. 55 p. (Series: The Study, vol. 2;

2. (b) **REPORTS**
 Main collection of articles is intended for scientific research workers
 concerned with problems of protection of health at nuclear powerplants. The volume contains 43 papers
 presented by Soviet scientists at the Second Conference on Nuclear Basis of
 Acute Mortality, held in Odessa in September 1970.

2. (b) **REPORTS**
 It is divided into two parts. Part I contains 27 papers dealing with
 problems on nuclear powerplants, facilities problems of particle emulsion and of
 dust removal. The first part is divided into two sections: "problems of protection of health at
 nuclear powerplants and control of their nuclear accidents".

Part II deals with particular problems in each field. The
 papers in Part II deal in detail with various problems in nuclear powerplants
 and in the choice of heavy stone and shale leaching, with the study of
 the mechanism of artificial earth walls and roads, described
 in a paper by A.E. Permyakov. The main theme of the presentation of
 the conference is published in the volume. The first 6 volume consists of 13
 reports presented by Soviet scientists on nuclear safety. Volume (1), Radiation
 Protection (Part 1); Volume (2), Radiation Protection (Part 2); Volume (3), Radiation
 Protection and Radiation Physics; Volume (4), Radiation Protection and
 Radiation Physics (Part 1); Volume (5), Radiation Protection and Radiation
 Physics (Part 2); Volume (6), Radiation Protection and Radiation Physics
 (Part 3); Volume (7), Radiation Protection and Radiation Physics (Part 4); Volume
 (8), Radiation Protection and Radiation Physics (Part 5); Volume (9), Radiation
 Protection and Radiation Physics (Part 6); Volume (10), Radiation
 Protection and Radiation Physics (Part 7); Volume (11), Radiation
 Protection and Radiation Physics (Part 8); Volume (12), Radiation
 Protection and Radiation Physics (Part 9); Volume (13), Radiation
 Protection and Radiation Physics (Part 10).

2. (b) **REPORTS**
 The second part of the collection is intended for non-medical scientists. In the present volume
 there are 16 articles which were written by Soviet scientists. The present
 work is aimed at those articles which were not identical

with the article by V.V. Gerasimov, V.N. Savchenko, V.A. Savchenko, et al., "The
 Chemistry of Flame Oxidation", and Polyakov, "Investigation of the
 Flame". The serial number of reports 230 and 236 are reported in the
 second edition.

2. (b) **REPORTS**
 Report on Soviet Scientists' Studies (1960-1965)
 Savchenko, Dr. on Nuclear Power by V. Savchenko (Report 207)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.D. Savchenko, Z.B. Savchenko, and
 S.Ye. Savchenko (Report 208)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.P. Savchenko, and Z.B. Savchenko (Report 209)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.P. Savchenko, and Z.B. Savchenko (Report 210)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.P. Savchenko, and Z.B. Savchenko (Report 211)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.P. Savchenko, and Z.B. Savchenko (Report 212)

2. (b) **REPORTS**
 Savchenko, Dr. A.A. Savchenko, L.P. Savchenko, and Z.B. Savchenko (Report 213)

2. (b) **REPORTS**
 Institute of Chemical Physics (VFFI) of the USSR Academy of Sciences (1970-1975)

21(1), 24(5)

AUTHOR:

Zaretskiy, D. F.

SOV/56-36-3-54/71

TITLE: On the Theory of Nuclear Magnetic Moments
(K teorii magnitnykh momentov yader)PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki,
1959, Vol 36, Nr 3, pp 869-873 (USSR)ABSTRACT: The important part played by spin-orbit coupling in the theory
of nuclear shells is, in general, taken into account by (1) :

$$\mathcal{X}_{ls} = -(\lambda \hbar/m^2c^2) \vec{s} [\nabla U \vec{p}]$$

(λ = phenomenological spin-orbit coupling constant,
 \vec{s} = nucleon spin operator, $U(\vec{r})$ = selfconsistent nuclear
 potential, \vec{p} = momentum operator of the nucleon); if the
 nucleus is in an electromagnetic field, (1) with $\vec{p} \rightarrow \vec{p} + e\vec{a}/c$ goes
 over into

$$\mathcal{X}' = (\lambda e \hbar/m^2c^3) \vec{s} [\nabla U \vec{a}] \quad (2)$$

where a is the vector potential. Mayer and Jensen (Ref 1)
 were the first to point out the necessity of taking this
 spin-orbit effect into account without, however, using (2).

Card 1/5

ON THE THEORY OF NUCLEAR MAGNETIC MOMENTS

SOV/56-56-3-54/71

D. I. Gorchukhin (Ref 2) calculated the spin-orbit coupling for magnetic transitions of the kind $d_{3/2} \rightarrow s_{1/2}$. In the present paper the effect is investigated by which (2) is introduced into the magnetic moment of spherical and non-spherical nuclei with odd proton number. Although (2) is not the only cause of the deviation of the observed magnetic moments from the Schmidt (Shmidt) line, an investigation (estimate) of (2) is nevertheless of interest, because it facilitates separation of the other effects. In some cases (2) makes the main contribution to the deviation of magnetic moments from the Schmidt line as well as to the probability of forbidden M1-transitions for nuclei with an odd proton number. In the case of nuclei with odd neutron numbers, consideration of (2) is of good purpose when there exists an effective charge due to interaction between neutron and the core of the nucleus (cf. Ref 3). In the present paper the additional factor occurring for strong spin-orbit coupling by the interaction between the nucleons in the nucleus and the electromagnetic field is, essentially, calculated for spherical and nonspherical nuclei. For the former the magnetic moment is $\mu = \langle J, J | \hat{\mu}_z | J, J \rangle$

Card 2/3

On the Theory of Nuclear Magnetic Moments

SOV/56-36-3-34/71

and, in consideration of (2):

$$\mu = \frac{1}{\pi} \frac{J + \frac{1}{2}}{2(J+1)} \frac{e \lambda}{2mc} \frac{\lambda}{mc^2} \langle J | r \frac{\partial U}{\partial r} | J \rangle$$

For Ru^{209} an estimate results in $\mu \approx +0.6 e \lambda / 2mc$.For nonspherical nuclei it holds that $\mu = \langle \hat{\mu}_J^2 \rangle / (J + 1)$

and with (2) for spin 1/2 also a formula, (21), is derived.

For Tm^{169} this results in $\mu = -0.35$ nuclear magnetons, as
against an experimental value of $\mu = -0.20 \pm 0.5$.
The author finally thanks P. E. Nemirovskiy for discussions.
There are 13 references, 7 of which are Soviet.

SUBMITTED: September 16, 1958

Card 3/3

24(5)
AUTHOR:

Zaretskiy, D. F.

SOV/56-36-4-26/70

TITLE:

On the Radiative Transitions in the System of
Rotational Levels of Nuclei With Spin 1/2 (O
radiatsionnykh perekhodakh v sisteme vrashchatel'nykh
urovney yader so spinom 1/2)

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1959,
Vol 36, Nr 4, pp 1129-1132 (USSR)

ABSTRACT:

For the purpose of calculating the probabilities of
magnetic dipole- and electric quadrupole transitions
between rotational levels of deformed spin 1/2 nuclei, a
coupling system suggested by Hund and already previously
used by the author (Ref 1) is used. A transformation of
coordinates (transition to the system of coordinates of
the nuclear axes) is carried out in the wave function,

$$\text{and by using the relation } \chi_{J_1, m_s} = \sum_{m_s'} D_{m_s m_s'}^{J_1} \chi_{J_1, m_s'}$$

Card 1/3

On the Radiative Transitions in the System of
Rotational Levels of Nuclei With Spin 1/2

SOV/56-36-4-26/70

a wave function of the form

$$\Psi^{JM} = \sqrt{(2J+1)/8\pi^2} \psi_0 \sum_{m_s} (\chi^{1/2} 0 m_s | \chi^{1/2} J m_s) D_{M, m_s}^J \chi_{J, m_s}$$

is obtained, which causes no difficulties with respect to calculating the probabilities of electric quadrupole transitions. The result agrees with that obtained by Bohr and Mottelson (Ref 2). (ψ_0 denotes the wave function describing the motion of the nucleon with respect to the nuclear axes, χ - the spin function of an odd nucleon in the laboratory system, the rotation quantum number χ assumes the values 0, 2, 4, ... for the states \sum_g^+ and 1, 3, 5, ... for \sum_u^+ . In the following the authors calculate the probabilities of magnetic dipole transitions and show that transitions between components of different doublets with $\Delta J = 1$ occur mainly as magnetic dipole transitions. The admixture of electric quadrupole transitions amounts to not more than 10 - 20 %. In

Card 2/3

On the Radiative Transitions in the System of
Rotational Levels of Nuclei With Spin 1/2

sov/56-36-4-26/70

conclusion, the nucleus Tm¹⁶⁹ (and Tm¹⁷¹) is shortly investigated as an example, and a comparison is drawn with the results obtained by references 7 and 8 with respect to the nucleus Yb¹⁷¹, the daughter nucleus originating from the β -decay of Tm¹⁷¹ (Fig). The author thanks S. A. Baranov for discussing experimental data, and D. P. Grechukhin for his comments. There are 1 figure and 9 references, 3 of which are Soviet.

SUBMITTED: September 13, 1958

Card 3/3

S
ZARETZKIY, D.F.

Radiative capture of a neutron. Zhur.eksp.i teor.fiz. 37 no.4:
1084-1087 0 '59. (MIRA 13:5)
(Neutrons--Capture)

16.8100, 24.6720, 24.6800, 24.6810

77009
SOV/56-37-6-49/55

AUTHORS: Zaretskiy, D. F., and Novikov, V. M.

TITLE: Letter to the Editor. Depolarization of Muons in
 μ -Mesoatoms with Deformed Nuclei

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki,
1959, Vol 37, Nr 6, pp 1824-1825 (USSR)

ABSTRACT: An analysis was made of the additional depolarization caused by the interaction of muon with nuclear deformation in μ -mesoatoms. The effect was considered for even-even nuclei. The Hamiltonian of the muon-nucleus system was taken in the following form:

$$H = H_0 + H_R + H_q$$

Here, H_0 is the Hamiltonian of the muon in the monopole field; H_R is operator of rotational energy of the

Card 1/4

Letter to the Editor. Depolarization of
Muons in μ -Mesocatoms with Deformed
Nuclei

77009
SOV/56-37-6-49/55

nucleus; H_q is operator of quadrupole interaction of muon with nucleus. The diagonalization of the Hamiltonian shows that the quadrupole interaction considerably changes the eigen functions of the system with corresponding muon in the 2p-states (cf., L. Wilets, Kong. Dansk. Vidensk. Selsk. Mat.-fys. Medd., 29, 3, 1954). The changes of the eigen functions of other states can be neglected. In such an approximation the polarization of muon in 1s-states becomes:

$$P = A_q W_{1/2} \langle \sigma_{2p_{1/2}} \rangle_0 + B_q W_{3/2} \langle \sigma_{2p_{3/2}} \rangle_0$$

Here $W_{1/2}$ and $W_{3/2}$ are probabilities of the passing of muon through states $2p_{1/2}$ and $2p_{3/2}$; $\langle \sigma_{2p_{1/2}} \rangle_0$ and

Card 2/4

Letter to the Editor. Depolarization of
Muons in μ -Mesocatoms with Deformed
Nuclei

77009
SOV/56-37-6-49/55

$\langle \sigma_{2p_{3/2}} \rangle_0$ are polarization of muon in the above states in the absence of quadrupole interaction; A_q and B_q are factors compensative of the additional depolarization. The factor B_q was determined from muon transition from the upper levels into the 1s-state through the states that can be described by the eigen function of the above given Hamiltonian:

$$B_q = 0.8 \sum_{k=1}^3 C_k \left[\left(\frac{E_{f.s.} - 3E_h}{3E_q} \right)^2 + 2 \right] - 0.6,$$

where

$$C_k^2 = \frac{E_q^2 (E_R + E_q - \frac{2}{3} E_{f.s.} - E_h)^2}{E_q^2 (E_R + E_q - \frac{2}{3} E_{f.s.} - E_h)^2 + E_q^2 (E_q + \frac{1}{3} E_{f.s.} - E_h)^2 + [(\frac{1}{3} E_{f.s.} - E_h) (E_R - \frac{2}{3} E_{f.s.} - E_h) - E_q^2]^2} \rightarrow \\ \rightarrow \frac{E_q^2 (E_R + E_q - \frac{2}{3} E_{f.s.} - E_h)^2}{- E_h (E_R - \frac{2}{3} E_{f.s.} - E_h) - E_q^2}.$$

Card 3/4

Letter to the Editor. Depolarization of
Muons in μ -Mesoatoms with Deformed

77009
SOV/56-37-6-49/55

Here, E_k is eigen values of the Hamiltonian which correspond to muon in 2p-states and to the whole momentum of the system 3/2; $E_{f.s.}$ is splitting of fine structure of the 2p-level; E_R is energy of the first rotational level of the nucleus; $E_q = \langle 2p | H_q | 2p \rangle$. The following values of B_q and D_q were calculated for Gd^{158} , W^{184} , Th^{232} , and U^{238} , respectively: 0.63, 0.64, 0.38, 0.38 and 0.61, 0.64, 0.6, 0.59. Thus, the interaction of muon with nuclear quadrupole deformation can lead to a considerable additional depolarization of the muon. There are 4 references; 2 Soviet, 1 Dutch, 1 U.S. The U.S. reference is: G. W. Ford, C. J. Mullin, Phys. Rev., 108, 477, 1957.

SUBMITTED: August 21, 1959

Card 4/4

GRIN', Yu.T.; DROZOV, S.I.; ZARETSKIY, D.F.

Green's function for odd nuclei. Zhur. eksp. i teor. fiz. 38
no.1:222-228 Jan '60. (MIRA 14:9)
(Potential, Theory of) (Nuclei, Atomic)

27480
S/048/61/025/009/005/007
B104/B102

24.6300

AUTHORS: Grin', Yu. T., and Zaretskiy, D. F.

TITLE: Collective excitations of non-spherical nuclei

PERIODICAL: Akademiya nauk SSSR. Izvestiya. Seriya fizicheskaya, v. 25,
no. 9, 1961, 1169 - 1175

TEXT: This paper was read at the 9th Annual Conference on Nuclear Spectroscopy. The authors generalize the theory of collective excitations of non-spherical nuclei. An equation is set up for the frequencies of collective nuclear oscillations, and a relation between the frequencies of the excited $\beta(k=0)$ and $\gamma(k=2)$ vibrational levels is derived in quasi-classical approximation. Using the Green's two-particle function

$K = \langle \Phi_0 | T(a_1 a_2^\dagger a_3 a_4^\dagger) | \Phi_0 \rangle$, the equation

$$1 = \chi \sum_{12} \frac{(E_1 E_1 - \epsilon_1 \epsilon_2 + \Delta^2)}{2E_1 E_2 [\omega^2 - (E_1 + E_2)^2]} |(q_{2k})_{12}|^2 \quad (9)$$

Card 1/5

27400

S/048/61/025/C09/005/007
B104/B102

Collective excitations of non-

is obtained for the frequencies of the bound states of two quasi-particles. Within the framework of the Nilsson model Eq. (9) can only be solved numerically for a real model. Using the model of an axisymmetrically deformed oscillator potential, the authors attempt to find a solution to this equation. Eq. (9) is represented in the form

$$1 = \chi \sum_{\lambda\lambda'} \frac{2\Delta^2 |q_{\lambda\lambda'}|^2}{E_\lambda (4E_\lambda^2 - \omega^2)} + \chi' \sum_{\lambda\lambda'}' \frac{(E_\lambda E_{\lambda'} - e_\lambda e_{\lambda'} - \Delta^2)(E_\lambda + E_{\lambda'})}{2E_\lambda E_{\lambda'} [(E_\lambda + E_{\lambda'})^2 - \omega^2]} |q_{\lambda\lambda'}|^2, \quad (15)$$

The first term is the sum over all transitions without energy change, and the second term is the sum over all other transitions. Since ω is negligible in the latter sum, Eq. (15) can be reduced to

$$1 = \chi' \sum_{\lambda\lambda'} \frac{2\Delta^2}{E_\lambda (4E_\lambda^2 - \omega^2)} |q_{\lambda\lambda'}|^2, \quad (16)$$

where

$$\chi' = \frac{\chi}{1 - \chi \sum_{\lambda\lambda'} \frac{E_\lambda E_{\lambda'} - e_\lambda e_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'} (E_\lambda + E_{\lambda'})} |q_{\lambda\lambda'}|^2}$$

Card 2/5

Collective excitations of non- ...

²⁷⁴⁸⁰
S/048/61/025/009/005/007
B104/B102

In quasi-classical approximation, Eq. (16) leads to

$$1 = \frac{x'}{\omega^2 \sqrt{\frac{4\Delta^2}{\omega^2} - 1}} \arctan \frac{1}{\sqrt{\frac{4\Delta^2}{\omega^2} - 1}} \sum_{\lambda\lambda'} |q_{\lambda\lambda'}|^2 \delta(\epsilon_{\lambda}) \quad \omega \sim \epsilon_0 / AR_0^4.$$

The following explicit solutions are obtained from this equation:

$$\omega = 2\Delta \sqrt{\frac{3}{2}} \cdot \sqrt{1 - x'} \sum_{\lambda\lambda'} |q_{\lambda\lambda'}|^2 \delta(\epsilon_{\lambda}) \quad \text{for } \omega \ll 2\Delta \text{ and}$$

$$\omega = 2\Delta \left\{ 1 - \frac{5}{8} x'^2 \left[\sum_{\lambda\lambda'} |q_{\lambda\lambda'}|^2 \delta(\epsilon_{\lambda}) \right]^2 \right\} \quad \text{for } 2\Delta = \omega \ll 2\Delta.$$

An analysis of these solutions indicates that collective excitations with energies much less than 2Δ may exist in deformed nuclei. β^- and γ^- vibrations are interrelated by

Card 3/5

27480
S/048/61/025/009/005/007
B104/B102

Collective excitations of non-

$$\frac{\omega_\beta^2}{\omega_\gamma^2} = \frac{\sum_{\lambda\lambda'} \frac{1 - g \left(\frac{e_\lambda - e_{\lambda'}}{2\Delta} \right)}{\left(\frac{e_\lambda - e_{\lambda'}}{2\Delta} \right)^3} (\eta_{22})_{\lambda\lambda'}^2 \delta(e_\lambda)}{\sum_{\lambda\lambda'} \frac{1 - g \left(\frac{e_\lambda - e_{\lambda'}}{2\Delta} \right)}{\left(\frac{e_\lambda - e_{\lambda'}}{2\Delta} \right)^3} (\eta_{40})_{\lambda\lambda'}^2 \delta(e_\lambda)} \quad (23)$$

From this relation, it is concluded that the frequencies of β - and γ -vibrations agree within the framework of the model of an axisymmetric oscillator potential. The agreement of the frequencies is closely related to the degeneracy of the levels in the oscillator potential. Finally, the Nilsson model without spin-orbital coupling is briefly discussed. The following values are obtained for the ratio $\omega_\beta/\omega_\gamma$ as a function of $\Delta = 2D\varepsilon_0/\omega_0$:

Δ	1.8	2.25	3.0	4.5
$\omega_\beta/\omega_\gamma$	0.90	0.86	0.80	0.72

S. T. Belyayev (Zh. eksperim. i teor. fiz. 39, 1387 (1960)) is mentioned.
Card 4/5

Collective excitations of non- ...

27480
8/048/61/025/009/005/007
B104/B102

There are 11 references: 7 Soviet and 4 non-Soviet. The references to English-language publications read as follows: Marumori T., Progr. Theor. Phys., 24, 331 (1960); Baranger M., Phys. Rev., 120, 957 (1960); Perlman I., Proceedings of the International Conference on Nuclear Structure, p. 547, Kingston, Canada, 1960.

Card 5/5

DROZDOV, S.I.; ZARETSKIY, D.F.

Effects of pair correlation near closed shells. Zhur. eksp. i
teor. fiz. 40 no.1:286-295 Ja '61. (MIRA 14:6)
(Potential, Theory of) (Superconductivity)

22151

S/056/61/040/003/031/031
B112/B214*24.6900*AUTHORS: Zaretskiy, D. F., Novikov, V. M.TITLE: Nuclear fission by μ -mesonsPERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 40,
no. 3, 1961, 982-983

TEXT: The effect of a meson on nuclear fission can be calculated for the case of rigidly oriented nuclear axes. The energy E_μ of a bound meson depends on the deformation parameters of the nucleus. A solution of Schrödinger's equation for a meson in the Coulomb field of the deformed nucleus is required for the determination of E_μ . It is assumed in the present paper that the nucleus has the form of an ellipsoid of rotation up to its saddle point. The Coulomb potential of a homogeneously charged ellipsoid of rotation with the semiaxes a and b has the form:

Card 1/3

22151

S/056/61/040/003/031/031

B112/B214

~~X~~

Nuclear fission by...

$$\varphi(a, \beta) = \frac{Ze}{c} \left\{ [1 - P_2(\text{ch } \alpha) P_2(\cos \beta)] \ln \text{cth} \frac{\alpha_0}{2} + \frac{3}{2} \frac{\text{ch}^2 \alpha}{\text{ch} \alpha_0} P_2(\cos \beta) + \right.$$

$$\left. + \frac{3}{4} \left(1 - \frac{\text{sh}^2 \alpha}{\text{sh}^2 \alpha_0} \right) \frac{\sin^2 \beta}{\text{ch} \alpha_0} \right\} \quad \text{for } \text{ch } \alpha \leq \text{ch } \alpha_0 = \frac{a}{c},$$

$$\varphi(a, \beta) = \frac{Ze}{c} \left\{ [1 - P_2(\text{ch } \alpha) P_2(\cos \beta)] \ln \text{cth} \frac{a}{2} + \frac{3}{2} \text{ch } \alpha P_2(\cos \beta) \right\} \quad (I)$$

for $\text{ch } \alpha \geq \frac{a}{c}$,

Here, Ze is the nuclear charge, $c^2 = a^2 - b^2$, P is the Legendre polynomial, and α, β are the degenerate elliptical coordinates. The Schrödinger equation for a bound meson in a nuclear field with such a potential was solved numerically with the help of an electronic computer.

The value of $E_{\mu}^{\text{thresh.}}$ for different values of a/b is given in the

following table (U^{238} nucleus):

a/b	1.2	1.4	1.6	1.8	2	2.2	2.5
$E_{\mu}^{\text{thresh.}} (\text{Mev})$	11.89	11.78	11.66	11.53	11.36	11.21	11.01

The increase ΔE of the fission barrier for some nuclei is given in another table:

Card 2/3

2215I

S/056/61/040/003/031/031
B112/B214

Nuclear fission by...

Nucleus	a/b statistical	a/b saddle point	E _{threshold} , Mev	ΔE, Mev
U ²³⁸	1.30	2.24	5.8	0.6
U ²³⁵	1.25	2.2	5.75	0.6
Pu ²³⁹	1.30	2.17	5.48	0.5

D. P. Grechukhin is thanked for his advice and V. K. Saul'yev for setting up the program and calculation with the electronic computer. There are 2 tables and 7 references: 2 Soviet-bloc and 3 non-Soviet-bloc.

SUBMITTED: January 11, 1961

Card 3/3

ZARETSKIY, D.F.; NOVIKOV, V.M.

Excitation of nuclei in heavy μ -mesic atoms. Zhur.ekspl teor.fiz.
41 no.1:214-221 J1 '61. (MIRA 14:7)
(Nuclei, Atomic)

ZARETSKIY, D.F.; URIN, M.G.

Nature of collective levels in nonspherical nuclei. Zhur.eksp.
i teor.fiz. 41 no.3:898-906 S '61. (MIRA 14:10)

1. Moskovskiy inzhenerno-fizicheskiy institut.
(Nuclei, Atomic)

ZARETSKIY, D.F.; URIN, M.G.

Microscopic description of collective levels of nonspherical nuclei.
Zhur. eksp. i teor. fiz. 43 no.3:1021-1030 '62. (MIRA 15:10)

1. Moskovskiy inzhenerno-fizicheskiy institut.
(Quantum theory) (Nuclie, Atomic)

S/056/62/042/001/045/048
B154/B112

AUTHORS: Zaretskiy, D. F., Urin, M. G.

TITLE: Low levels with negative parity in nonspherical nuclei

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 42,
no. 1, 1962, 304-305

TEXT: In deformed even-even nuclei, low levels with 0.3-1.3 Mev were recently detected (Ref. 1). These 1⁻ levels correspond to internal excitations with K = 0 and negative parity. They cannot be explained by the hydrodynamic model. Therefore, it was proposed that such levels should be interpreted in terms of the superfluid model. Using the method of approximate second quantization in the same way as in Ref. 1 (Zaretskiy, D. F., Urin, M. G., ZhETF, 41, 398, 1961), the excitation energy ($\hbar = 1$) is found to be

$$1 = \times \sum_{\lambda\lambda'} n_p |(q_{30})_{\lambda\lambda'}|^2 \frac{E_\lambda E_{\lambda'} - \epsilon_\lambda \epsilon_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'}} \frac{E_\lambda + E_{\lambda'}}{(E_\lambda + E_{\lambda'})^2 - \omega^2} \quad (1)$$

Card 1 / 5

S/056/62/042/001/046/048
B154/B112

Low levels with negative ...

where $q_{30} = r^3 \cdot Y_{30}(\theta)$ is the single operator for the octupole moment; $E_\lambda = \sqrt{\varepsilon_\lambda^2 + \Delta^2}$. The energy ε_λ is subtracted from the Fermi surface; $\Delta = \Delta_n(\Delta_p)$ is a constant characterizing the energy of proton (neutron) pair correlation; $\varepsilon_o^n = \varepsilon_o^n(\varepsilon_o^p)$ is the energy of the Fermi boundary for neutrons (protons); $\kappa \sim \varepsilon_o / AR^6$ is constant depending on the octupole-octupole interaction. The value of κ is proposed to be the same for nn, pp, and np interactions. (1) permits two solutions which, at $\kappa \rightarrow 0$, correspond to the dissociation of a neutron or a proton pair. For a given group of nuclei, κ can be regarded as constant. If κ is therefore determined for one group of nuclei from the position of the 1⁻ level, it will be possible to estimate ω from Eq. (1) for the other nuclei of this group. The probability of electric dipole transition from the 1⁻ level to the ground level is expressed by ✓

Card 2/5

Low levels with negative ...

S/056/62/042/001/046/048
B154/B112

$$B(E1; I^- \rightarrow 0^+) = \frac{1}{6\omega} \left\{ \sum_{\lambda, \lambda'} e_{n, p} (q_{10})_{\lambda \lambda'} (q_{30})_{\lambda \lambda'} \frac{E_\lambda E_{\lambda'} - e_\lambda e_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'}} \frac{E_\lambda + E_{\lambda'}}{(E_\lambda + E_{\lambda'})^2 - \omega^2} \right\}^2 \times \\ \times \left\{ \sum_{\lambda, \lambda'} |(q_{30})_{\lambda \lambda'}|^2 \frac{E_\lambda E_{\lambda'} - e_\lambda e_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'}} \frac{E_\lambda + E_{\lambda'}}{|(E_\lambda + E_{\lambda'})^2 - \omega^2|^2} \right\}^{-1} \quad (2),$$

where $e_n = -Ze/\Lambda$, $e_p = Ne/\Lambda$, e is the proton charge, and $Z(N)$ is the number of protons (neutrons). In quasiclassical approximation, the authors estimate $B(E1)$ for low I^- levels ($\omega \ll 2\Delta_{n, p}$) according to (Ref. 4):

$$\sum_{\lambda, \lambda'} |(q_{30})_{\lambda \lambda'}|^2 \frac{E_\lambda E_{\lambda'} - e_\lambda e_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'} (E_\lambda + E_{\lambda'})^2} \approx \quad (3), \\ \approx \frac{1}{4\Delta^2} \sum_{\lambda, \lambda'} |(q_{30})_{\lambda \lambda'}|^2 \varphi \left(\frac{e_\lambda - e_{\lambda'}}{2\Delta} \right) \delta(e_\lambda) \sim \rho_0 R^6 \Delta^{-2},$$

where $\varphi(x) = x^{-2} \ln(x + \sqrt{1 + x^2}) \cdot [x^3 \sqrt{1 + x^2}]^{-1}$, and ρ_0 is the level energy density.

Card 3/5

S/056/62/042/001/046/048
B154/B112

Low levels with negative ...

density near the Fermi surface. For the approximation in (3), $\epsilon_\lambda - \epsilon_{\lambda'} \lesssim 2\Delta$ is necessary. From the Nilsson scheme it follows that in the observed nuclear quadrupole deformations β_0 there are levels λ, λ' for which this condition is satisfied. Analogous evaluation of the numerator in Eq. (2) leads to

$$B(E1, 1^- \rightarrow 0^+) \sim \left(\frac{N-Z}{A} \right)^2 (eR)^2 \beta_0^2 \rho_0 \Delta \frac{2\Delta}{\omega} \quad (4).$$

In the model considered, the probability for the excitation of the 3^- level is

$$B(E3, 0^+ \rightarrow 3^-) = \frac{1}{2\omega} \left\{ \sum_{\lambda\lambda'} p |(q_{30})_{\lambda\lambda'}|^2 \frac{E_\lambda E_{\lambda'} - \epsilon_\lambda \epsilon_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'}} \frac{E_\lambda + E_{\lambda'}}{(E_\lambda + E_{\lambda'})^2 - \omega^2} \right\}^{-1} \times \quad (5). \\ \times \left\{ \sum_{\lambda\lambda' p} |(q_{30})_{\lambda\lambda'}|^2 \frac{E_\lambda E_{\lambda'} - \epsilon_\lambda \epsilon_{\lambda'} + \Delta^2}{2E_\lambda E_{\lambda'}} \frac{E_\lambda + E_{\lambda'}}{|(E_\lambda + E_{\lambda'})^2 - \omega^2|^2} \right\}^{-1}.$$

This relation is obtained without introduction of the effective nucleonic charge for the $E3$ transition. Estimating $B(E3, 0^+ \rightarrow 3^-)$ in analogy to (3)
Card 4/5

S/056/62/042/002/032/055
B108/B104

AUTHORS: Zaretskiy, D. F., Novikov, V. M.

TITLE: Excitation of collective nuclear levels in heavy μ -mesic atoms

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 42,
no. 2, 1962, 511 - 519

TEXT: The excitation probability of low collective levels with spin 2^+ which depends on the magnitude of the nuclear quadrupole moment and on the quadrupole transition probability for the level in question is discussed for heavy mesic atoms. Not only a passing $2p$ muon but also a $3d$ muon may excite the rotational levels of heavy Th or U type nuclei. The interaction part of the Hamiltonian of the nucleus-muon system is ascribed to quadrupole interaction of the levels $2p_{1/2}$ and $2p_{3/2}$ only. Results of the calculations given in Table 1 indicate that the excitation probability W depends on the nature of the collective level. W is increased, however, by some 20% owing to $3d_{3/2}$ and $3d_{5/2}$ muon states. The latter can

✓

Card 1/3

Excitation of collective nuclear...

S/056/62/042/002/032/055
B108/B104

also lead to the excitation of the 4^+ level. The excitation probability W_{4+} , however, is much more sensitive to the sign of the quadrupole moment than is W_{2+} . The hyperfine splitting of the excited nuclear levels owing to the interaction between the magnetic moments of the muon and of the excited nucleus is calculated. The finite size of the nucleus has to be considered. For Th²³² and U²³⁸, $\Delta E = 840$ ev. The additional depolarization of muons owing to the transfer of polarity to a nucleus with excited collective levels is also discussed. However, this effect may be masked by the effect of the electron shell. The circular polarization of gammas from the $2p - 1s$ transition may give information about the additional depolarization of the muons. The effects of quadrupole interaction may also have an effect upon the shape of the muon transition lines. In general, the effect of quadrupole excitation of collective levels in heavy mesic atoms may be masked by radiationless excitation. The latter effect may be estimated by studying the nuclear transitions $4^+ \rightarrow 2^+$ and $2^+ \rightarrow 0^+$ with respect to the occurring gamma quanta from the muon transition. There are 1 figure, 4 tables, and 13 references: 8 Soviet and 5 non-Soviet.

Card 2/3

Excitation of collective nuclear...

S/056/62/042/002/032/055
B108/B104

The four references to English-language publications read as follows:
 Jacobson. Phys. Rev., 96, 1637, 1954; M. V. Patter. Nucl. Phys., 14, 42, APPROVED FOR RELEASE 09/19/2001 M. V. Patter. Nucl. Phys., 11, 419, 1959; A. Z. Dolginov. Nucl. Phys., 7, 569, 1958.

SUBMITTED: August 8, 1961

Table 1. Results.

Legend: (1) nucleus; (2) parameter of axial symmetry; (3) spin-orbital splitting; (4) energy of the 2^+ level; (5) excitation probability.

(1)	$B(E2) \cdot 10^4$ eV	(2) Y_{spad}	$ Q_{M1} ^2$ $B(E2)$	(3) ΔE MeV	(4) E_{2+} MeV	(5) W_{2+}
W ¹⁸⁸	356	15,8	3,86	0,134	0,123	0,40*
O ₃ ¹⁸⁰	255	22,3	2,68	0,147	0,187	0,30
Hg ¹⁹⁸	413	24,3	1,80	0,162	0,411	0,025

Card 3/3

40429

S/056/62/043/003/043/063
B108/B102

24.4400

AUTHORS: Zaretskiy, D. F., Urin, M. G.

TITLE: Microscopic description of collective levels of nonspherical nuclei

PERIODICAL: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 43,
no. 3(9), 1962, 1021 - 1030

TEXT: When pair correlations of nucleons are considered by the method of Bogolyubov's canonical transformation this leads to unphysical states. Here the structure of such collective states in deformed nuclei is studied by second quantization approximation (cf. ZhETF, 41, 898, 1961). In this case the Hamiltonian for the collective excitations is represented in a diagonalized form wherefrom the collective unphysical state can be eliminated. The unphysical states of the form $A_1^+ |C_0\rangle$ are orthogonal to all physical (real) states. The error which results from simplifying the Hamiltonian $H = H_0 + H_Q + H_{int}$ due to collective oscillations is of the order of $(\epsilon_0 \Delta)^{-1} \ll 1$ where Δ is a constant determining the pairing energy.

Card 1/3

Microscopic description of...

S/056/62/043/003/043/063
B108/B102

and ρ_0 is the energy density of the single-nucleon levels near the Fermi level. The matrix element of the electrical multipole moment for a transition between ground state and excited state with multipolarity λ and projection K of the moment has the form.

$$\langle \lambda K | Q_{\lambda K}^{\ell} | 0 \rangle^2 = \left[e \sum_{1,2} |(q_{\lambda K})_{12}|^2 I_{12}(\omega_{\lambda K}) \right]^2 \left\{ \omega_{\lambda K} \sum_{1,2} |(q_{\lambda K})_{12}|^2 J_{12}(\omega_{\lambda K}) \right\}^{-1}, \quad (24)$$

with $J_{12}(\omega^2) = \partial I_{12} / \partial \omega^2$; $I_{12}(\omega_{\lambda K}) = (E_1 E_2 - \epsilon_1 \epsilon_2 + \Delta^2) E_{12} / 2 E_1 E_2 (E_{12}^2 - \omega_{\lambda K}^2)$; $E_v = \sqrt{\epsilon_v^2 + \Delta^2}$, $E_{12} = E_1 + E_2$, ϵ_v is the single-nucleon state energy. In the case of β -excitations, Eq. (24) determines the reduced $E\lambda$ -transition probability $B(E\lambda) \sim B_{sp}(E\lambda) \rho_0 \Delta^2 / \omega_{\lambda K}$ with an accuracy of $\sim \beta_0^2$ where β_0 is the equilibrium deformation of the nucleus and B_{sp} is the reduced single-particle transition probability.

Card 2/3

ZARETSKIY, D. F.; LUSHINKOV, A. A.

"Applications of the Theory of the Fermi Liquid to Nuclear Photoabsorption."

report submitted for All-Union on Nuclear Spectroscopy, Tbilisi, 14-22 Feb
64.

IAE (Inst Atomic Energy)

MIGDAL, A. B.; LUSHNIKOV, A. A.; ZARETSKIY, D. F.

"Theory of nuclear dipole photoabsorption."

report submitted for Intl Conf on Low & Medium Energies Nuclear Physics,
Paris, 2-8 Jul 64.

Kurchatov Inst, Moscow.

ZARETSKIY, D.F.; URIN, M.G.

Oscillations of nonspherical nuclei. Izv. AN SSSR. Ser. fiz. 28
no.1:118-126 Ja '64. (MIRA 17:1)

SOURCE: Zhurnal eksperimental'noy i teoreticheskoy fiziki, v. 48, no. 1, 1965,

ABSTRACT: The conditions are investigated, under which a gamma quantum radiated by one of the nuclei without recoil (Mossbauer effect) can be absorbed by another nucleus of the same type, so that the excitation (nuclear exciton) can propagate over an entire host crystal consisting of excited and unexcited nuclei of the same type. It is shown that when a nuclear exciton is produced, the main characteristics of spontaneous emission change markedly, because singular anisotropy appears and the gamma quantum flux is concentrated predominantly in the direction of the reciprocal-lattice vector. The emission probability in

this direction is

Card 1/2

L 24817-65

ACCESSION NR: AF5004413

estimated to be approximately equal to the square of the effective number of nuclei. It is also shown that the formation of a nuclear excision is connected with an increase in the width of the emitting level by a factor proportional to the cube root of the number of effective nuclei. It is thus possible to obtain sharply directional beams of monochromatic gamma quanta and the lifetimes of the nuclear isomers can be greatly reduced when such isomers are placed in a crystal consisting of unexcited nuclei of the same sort. "We thank Y. A. Shapiro for valuable remarks and also V. K. Vaytovetskiy and S. M. Feynberg for discussions." Orig. art. has: 30 formulas.

ASSOCIATION: Institut stekloj energii (Institute of Atomic Energy)

SUBMITTED: 28July4

EICL: 40

SUB CODE: 56, OP

IC REF ID: 014

OTHR: 003

Card 2/2

L 2752-66 EWT(m)/T/EWA(m)-2
ACCESSION NR: AF5024345

44,55

UR/0367/65/002/002/0307/0314

AUTHOR: Zaretskiy, D. F.; Ivanter, I. G.

44,55

46
B

TITLE: The four-fermion interaction and baryon masses

19 44,55

SOURCE: Yadernaya fizika, v. 2, no. 2, 1965, 307-314

TOPIC TAGS: particle symmetry, unitary symmetry, group theory, strong nuclear interaction, baryon, fermion, particle physics

ABSTRACT: A simple dynamic model is proposed for interpreting the mass spectrum of baryons (octet and decuplet). The mathematical analysis is based on a Hamiltonian with strong four-fermion interaction, which corresponds to a scalar and to the eighth component of an 8-vector $SU(3)$ representation. The four-fermion interaction is made up of strongly interacting baryon currents. This model is used as a basis for determining the relationship between splitting of the masses of the baryons in the octet and corresponding splitting in the decuplet. Orig. art. has: 32 formulas.

ASSOCIATION: none

SUBMITTED: 30Dec64

NO REF Sov: 002

MLR
Card 1/1

ENCL: 00

OTHER: 016

SUB CODE: HP, MA

SHAPIRO, I.S.; ZARETSKIJ, D.F.; LUSHNIKOV, A.A.

[Nuclear physics; the mechanism underlying nuclear reactions] IAdernaia fizika; mekhanizm iadernykh reaktsii.
Moskva, AN SSSR, 1965. 88 p. (MIRA 18:10)

1. Akademiya nauk SSSR. Institut nauchnoy informatsii.

L 36377-66 BMF(m)/T
ACC NR: AP6017590
SOURCE CODE: UR/0367/66/003/002/0263/0267
58
57
B

AUTHOR: Zaretskiy, D. F.; Lomonosov, V. V.
ORG: none
TITLE: Concerning certain features of the radiation of gamma quanta from nuclei in a crystal lattice 19
SOURCE: Yadernaya fizika, v. 3, no. 2, 1966, 263-267
TOPIC TAGS: gamma radiation, excited nucleus, nuclear isomer, nuclear energy level, crystal lattice structure, line width, spectral distribution, angular distribution
ABSTRACT: This is a continuation of earlier work by the authors (ZhETF v. 43, 368, 1965) where it is shown that under certain conditions collective nuclear excitation (nuclear exciton) can exist in an ideal single crystal, and that this exciton has a decay width which differs noticeably from the natural width of the nuclear level. The present article is devoted to a study of the singularities of radiative decay of nuclear levels in the case when the conditions for the existence of a nuclear exciton are present, and to a study of the spectral and angular distribution of the gamma quantum flux in this case. To this end, one of the methods of generation of nuclear collective excitation and the effects associated with it is investigated, and the radiation from a recoilless nucleus placed in a definite crystal lattice point is determined. All the nuclei in the crystal are assumed to be of the same kind as the radiating

Card 1/2

L 36371-66

ACC NR: AP6017590

nucleus, and line splitting is neglected. From the equations of motion the authors determine the amplitude of the state corresponding to the case when the isomer is excited but the other nuclei are not, the time dependence of the decay probability of the isomer, and the spectral and angular distribution of the gamma quanta emitted under definite relations between the wavelength of the radiation and the lattice constants. The results are extended to a three-level isomer and to radiation of more than one nucleus in the crystal. It is concluded that the existence of the nuclear exciton can be ascertained by observation of the spectral and angular distribution of the gamma-quantum flux when the condition $K = 2\pi b$ is satisfied (K - wave vector of the radiation, b - reciprocal lattice vector). The authors thank V. K. Vaytovetskiy for useful discussions. Orig. art. has: 16 formulas.

SUB CODE: 20/ SUBM DATE: 08Jan65/ ORIG REF: 002/ OTH REF: 001

me
Card 2/2

The effect of deformation on the corrosion of metals. E.M. Zarotskii. J. Applied Chem. U.S.S.R. 24, 521-8(1951)(Engl. translation).

See C.A. 46, 4984a.

B.R.

immediate source clipping


Influence of Deformation on the Corrosion of Metals. (In Russian.) E.N.
Zaretskii. Zhurnal Prikladnoi Khimii, v. 24, May 1951, p. 479-484.

See abstracts for parts I and II, Chemical Abstracts items 344-3 and 363-2,
1951.
(R general, R11, 424, Al, Zn, Cu, CP)

Immediate Source Clipping

ZARETSKII, E. N.

RT-1237 *A mechanism of corrosive cracking in magnesium alloys* / Abstracted from:
O mekhanizme korrozionnogo rastreskivaniia magnievykh splavov.
Doklady Akademii Nauk SSSR, 58 (4): 607-610, 1947



met. abstract

6

*Corrosive Cracking of Magnesium Alloys.—I. II. E. M.
Zaritsky. (Zhur. Tekhn. Khim., 1947, 20, 823-839, 830
and Brit. Abs., 1948, [R1], 493).—[In Russian]. [II.—]
Sheets of pure Mg and rolled alloy MA-1 (contg. 1.2% Mn)
do not exhibit flaking after exposure to atmospheric con-
ditions. MA-3 alloy (6.6% Al) develops cracks when exposed
to the weather under elastic stress, the time required for the
cracks to appear increasing from 1 day under a load of 16
kg./mm.² to 32-43 days under a load of 2.1 kg./mm.². These
cracks spread rapidly, passing through crystallites as well as
along their boundaries. Cracking cannot be prevented by
chem. or electrolytic oxidation of the surface, or by mech.
or thermal treatment of the alloy. [II.—] Cast Mg or MA-3
alloy sheets do not crack during weathering under elastic
stress. MA-6 alloy (10% Al) cracks readily under elastic
conditions, but this effect mostly disappears after annealing
at 420° C. for 3 hr., and quenching in H₂O.

*BJ-6, Non-ferrous metals**B1.16.*

Corrosive cracking of magnesium alloys. L. II. E. M. Zaretsky
(USSR Chem., USSR, 1947, 50, 533-529, 530-536). I. Sheris
of pure Mg and rolled electron alloy MA-1 (containing 1.22% of Mn)
do not exhibit flaking after exposure to atm. conditions. MA-3
alloy (5.8% of Al) develops cracks when exposed to the weather
under elastic stress, the time required for development of the cracks
rising from 1 day under a load of 16 kg. per sq. mm. to 32-43
days under a load of 8.1 kg. These cracks spread rapidly, passing
through crystallites as well as along their boundaries. Cracking
cannot be prevented by chemical or electrolytic oxidation of the
surface, or by mechanical or thermal treatment of the alloy.
II. Cast Mg or MA-3 alloy sheets do not crack during weathering
under elastic stress. MA-6 alloy (10% of Al) cracks readily under
these conditions, but this effect mostly disappears after annealing
at 420° for 3 hr., and quenching in H₂O. R. Tavscor.

BC

A-1

Corrosion of copper in aqueous ammonium salts. I. Mechanism of corrosion. E. M. Zanetikhan and G. V. Akutov (J. Appl. Chem. Russ., 1959, 11, 1161-1172).—Cu is not corroded by aq. NH_3 or NH_4Cl in absence of O_2 . In its presence corrosion consists of 3 successive processes: (i) dissolution of Cu, with formation of $\text{Cu}_2\text{O}-\text{NH}_3$ complexes (corrosion with O_2 depolarization), (ii) oxidation of Cu^{+} to Cu^{2+} , and (iii) electrochemical reaction of the Cu^{2+} complex with Cu, to regenerate the Cu^{+} complex. Reaction (iii) constitutes a separate cycle, and is clearly responsible for corrosion; during this part of the process the Cu becomes strongly electro-negative. The corrosive action of aq. NH_3 is > of NH_4Cl ; the velocity of reaction $\propto [\text{NH}_3]$ in the former case, and $\propto \text{max.}$ in aq. NH_4Cl ; in the latter, the relatively weaker action of NH_4Cl is ascribed to instability of the Cu complexes in its presence.
R. T.

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

SECOND DIVISION		SECTION MAY ONLY ONE		SECTION		SECOND DIVISION	
A	B	C	D	E	F	G	H

CA

9

The effect of deformation on the corrosion of metals.
E. M. Zaritskii. *J. Applied Chem. U.S.S.R.* 24, 821-8
(1951) (Engl. translation).—See C.A. 46, 10346. B.R.

MR

R

244-R. Influence of Deformation on
the Corrosion of Metals. (In Russian.)
E. M. Zaretskii. *Zhurnal Prikhodnoi
Khimii*, v. 24, May 1981, p. 477-484.
See abstracts of parts I and II,
Chemical Age; items 344-R and 345-
R, 1981.
(R general, RII, Q24, Al, Zn, Cu,
ST)

Int. Rev.
1952

P. - Physical Properties
and Test Methods

1952-P. The Influence of Deformation
on the Potentials of Metals. (In Russian.) E. M. Zaretskii. *Zhurnal Prik-
luden Khimii*, V. 25, June 1951, p. 614-
623.

See abstract of "Deformation and
Corrosion" (*Chemical Age*; Items 314-
B and 399-H, 1951)
(R11, Q24, Mg, Al, Cu, Zn, ST)

CIA

The mechanism of removal of oxides in the process of steel etching by acids. V. O. Krenig and E. M. Zaretskii. *Koroziya i Borba s Nej* 5, No. 3-4, 9-18 (1930). —

To prove the electrochemical nature of the process of etching, experiments were made with the aid of a microelectrode, one electrode of which was made of steel and the other of the sterilized sample having the general formula Fe_3O_4 . The steel, Cromanyl, contained C 0.4, Cr 1.01, Mn 1.01, Si 0.87, P 0.035 and Cu 0.10%. The oxides formed on this steel contain about 90.8% Fe_3O_4 + Fe_2O_3 , 1.4% of FeO and 1.2% of metallic Fe. As the process goes on the potential of the anode changes very little. During the first 10 min. a strong polarization takes place; then the potential of the cathode rises to a value slightly above the initial one; finally it drops again, almost reaching the value of the anode potential. Both the anode and the cathode undergo losses in weight. During the first two hrs. evolution of H was observed only on the anode. Then bubbles began to appear on the cathode accompanied by seepage of the oxide film from the Fe all along the cathode surface. By that time the current strength fell almost to zero. FeO becomes completely dissolved and metallic Fe decreases to 0.1%. After etching the magnetic properties of the oxide crust have almost disappeared, which means that Fe_3O_4 was largely reduced during the process, while Fe_2O_3 remained practically unchanged. The mechanism of etching can be represented as follows: On the anode Fe^{+2} ions are formed. On the microcathode regions of the anode the H^{+} ions become discharged. On the cathode the H^{+} ions

also become discharged but in the presence of the Fe_3O_4 they react to form Fe_2O_3 , which dissolves in the acid. The velocity of this reaction is less than that of the H-ion discharge. Therefore, soon H also begins to form and its accumulation causes the polarization of the cathode. Both the cathode potential and the current density drop until the velocities of the two processes become equal. With time the film of Fe_3O_4 disappears in some spots and the acid comes into contact with the underlying film of Fe_2O_3 , which has a higher tendency to oxidize. At that moment the cathode potential acquires a positive value, the potential difference and the v. d. increase and the discharged H ions reduce Fe_3O_4 to Fe_2O_3 , which goes into the acid. When most of the Fe_3O_4 is reduced the basis metal, i.e., Fe, becomes exposed to the action of the acid. The situation is the same now as on the anode. Thus on the micro-anode regions of the cathode H bubbles appear and the potential drops back to the same value as on the anode.

C. S. Shapiro

AIA-SEA METALLURGICAL LITERATURE CLASSIFICATION

SECTION SUBJECT		SECTION NO.		SECTION SUBJECT		SECTION NO.	
SOURCE NO.	SECTION NO.	SECTION NO.	SECTION NO.	SECTION NO.	SECTION NO.	SECTION NO.	SECTION NO.
1	2	3	4	5	6	7	8
9	10	11	12	13	14	15	16
17	18	19	20	21	22	23	24
25	26	27	28	29	30	31	32
33	34	35	36	37	38	39	40
41	42	43	44	45	46	47	48
51	52	53	54	55	56	57	58
61	62	63	64	65	66	67	68
71	72	73	74	75	76	77	78
81	82	83	84	85	86	87	88
91	92	93	94	95	96	97	98

CP

Comments on the paper of V. S. Zorin. Corrosion
process under drops of solutions. I. M. Vaynshteyn
J. Phys. Chem. (U.S.S.R.) 13, 1021-3 (1939). Cf. C.A.
33, 6213. Reply. V. S. Zorin. *J. Russ. Phys. Chem.*
V. H. Rathbun

APPENDIX 4 METALLURGICAL LITERATURE CLASSIFICATION


STANDARD SUBJECT	EXCERPT FILE CODE	SECTION	ALPHABETICAL INDEX
MATERIALS	M	1	A


C

Corrosion of copper in aqueous solutions of ammonium compounds. I. The mechanism of the corrosion. E. M. Zaretskii and G. V. Akimov. *J. Applied Chem.*

(U. S. S. R.) II, 1161-71 (in French, 1171-2) (1938).—The corrosion of Cu in solns. of NH₄ compds. is attributed to three cooperative processes: (1) electrochem. soln. of Cu with the formation of cuprous-ammonia complex, corresponding to a special type of corrosion process with the depolarization of O₂; (2) chem. oxidation of the complex formed by the O₂ of the air to cupric-ammonia complex; and (3) electrochem. reduction of the cupric-ammonia complex to cuprous-ammonia complex. The soln. of Cu proceeds energetically in process (3), which together with (2) forms an independent cycle. Because of inconsiderable dissoci. of the copper-ammonia complex cation, the Cu potential shifts strongly to the neg. side and the difference between anode and cathode reaches a considerable value. The Cu corrosion is less intensive in NH₄Cl soln., presumably because the complex is partially destroyed and therefore its potential is more pos. and the difference between the anode and cathode portions is smaller. In the absence of air, Cu is practically unaffected by corrosion in the NH₄OH and NH₄Cl solns. The velocity of Cu corrosion increases with the increase of NH₄OH concn.; it increases with NH₄Cl concn. to N and then decreases with further increase of the NH₄Cl concn. NH₄Cl soln. corrodes Cu to a smaller degree than does NH₄OH.

A. A. Podgorny







1ST AND 2ND COLUMNS		ACCESSION AND PREPARED BY INDEX		10D AND 11H COLUMNS	
<i>A</i>		664. Corrosion Cracking of Cast Magnesium Alloys. (In Russian). E. M. Zarataev. Zhurnal Prikladnoi Khimii (Journal of Applied Chemistry), v. 20, Sept. 1947, p. 830-840.		<i>R</i>	
		Stress-corrosion resistance of cast magnesium, of the alloy mentioned in the preceding abstract, and of another one containing 10% Al, 0.20% Mn, 0.04% Fe, and 0.09% Si, were investigated. The first was not sensitive to stress-corrosion; the cast form of the second, contrary to the cold worked form, is also not			
		Sensitive; the latter cast alloy has a slight tendency toward stress-corrosion, which can be eliminated by proper heat treatment and quenching procedure.			
COMMON ELEMENTS		MATERIALS INDEX		COMMON ELEMENTS	
ASME-SLA METALLURGICAL LITERATURE CLASSIFICATION		EXTRAS		EXTRAS	
11CHI STRAIN RATE		11CHI BONDING		11CHI BONDING	
STRIED #	STRIEDS H/M OR ONLY USE	COLLECTION	STRIED OR ONLY USE	STRIED OR ONLY USE	STRIED OR ONLY USE
1 2 3 4 5 6 7 8 9 10 11	1 2 3 4 5 6 7 8 9 10 11	1 2 3 4 5 6 7 8 9 10 11	1 2 3 4 5 6 7 8 9 10 11	1 2 3 4 5 6 7 8 9 10 11	1 2 3 4 5 6 7 8 9 10 11
12 13 14 15 16 17 18 19 20 21	12 13 14 15 16 17 18 19 20 21	12 13 14 15 16 17 18 19 20 21	12 13 14 15 16 17 18 19 20 21	12 13 14 15 16 17 18 19 20 21	12 13 14 15 16 17 18 19 20 21
22 23 24 25 26 27 28 29 30 31	22 23 24 25 26 27 28 29 30 31	22 23 24 25 26 27 28 29 30 31	22 23 24 25 26 27 28 29 30 31	22 23 24 25 26 27 28 29 30 31	22 23 24 25 26 27 28 29 30 31
32 33 34 35 36 37 38 39 40 41	32 33 34 35 36 37 38 39 40 41	32 33 34 35 36 37 38 39 40 41	32 33 34 35 36 37 38 39 40 41	32 33 34 35 36 37 38 39 40 41	32 33 34 35 36 37 38 39 40 41
42 43 44 45 46 47 48 49 50 51	42 43 44 45 46 47 48 49 50 51	42 43 44 45 46 47 48 49 50 51	42 43 44 45 46 47 48 49 50 51	42 43 44 45 46 47 48 49 50 51	42 43 44 45 46 47 48 49 50 51
52 53 54 55 56 57 58 59 60 61	52 53 54 55 56 57 58 59 60 61	52 53 54 55 56 57 58 59 60 61	52 53 54 55 56 57 58 59 60 61	52 53 54 55 56 57 58 59 60 61	52 53 54 55 56 57 58 59 60 61
62 63 64 65 66 67 68 69 70 71	62 63 64 65 66 67 68 69 70 71	62 63 64 65 66 67 68 69 70 71	62 63 64 65 66 67 68 69 70 71	62 63 64 65 66 67 68 69 70 71	62 63 64 65 66 67 68 69 70 71
72 73 74 75 76 77 78 79 80 81	72 73 74 75 76 77 78 79 80 81	72 73 74 75 76 77 78 79 80 81	72 73 74 75 76 77 78 79 80 81	72 73 74 75 76 77 78 79 80 81	72 73 74 75 76 77 78 79 80 81
82 83 84 85 86 87 88 89 90 91	82 83 84 85 86 87 88 89 90 91	82 83 84 85 86 87 88 89 90 91	82 83 84 85 86 87 88 89 90 91	82 83 84 85 86 87 88 89 90 91	82 83 84 85 86 87 88 89 90 91
92 93 94 95 96 97 98 99 100 101	92 93 94 95 96 97 98 99 100 101	92 93 94 95 96 97 98 99 100 101	92 93 94 95 96 97 98 99 100 101	92 93 94 95 96 97 98 99 100 101	92 93 94 95 96 97 98 99 100 101

AS M

248-R. Influence of Deformation on
the Corrosion of Metals. (In Russian.)
D. M. Zaretskii. *Zhurnal Prikladnoi
Khimii*, v. 24, May 1961, p. 477-484.
See abstracts of parts I and II,
Chemical Age; items 344-R and 368-
R, 1961.
(R general, R11, Q24, Al, Zn, Cu,
ST)





CA

9

Effect of deformation on the corrosion of metals. E. M. Lantsikil, Zhur. Priklad. Khim. (J. Applied Chem.) 24, 177-84(1951) -- Test strips of rolled and annealed Mg alloys with (I) 1.08% Mn, (II), Al 0.15, Ce 0.35, Mn 1.67%, (III) Al 5.62, Zn 0.06, Mn 0.33, Ni 0.17%, and 1 mm. thick steel C 0.10, Mn 0.32, Si 0.10, Ni 0.15, Cr traces; 0 mm. thick Cu; 3 mm. thick Zn; and 2 mm. thick Al were subjected to known tensile stresses σ , and the degree of corrosion was determined by the loss of wt. in 24 hrs. With I, stress up to 100% of the limit had no effect on the corrosion in a 0.1 M NaCl soln. (buffered to pH 6.8); in atm. exposure, corrosion first decreases with increasing σ , passes through a min. (at about $\sigma = 40\%$), then increases. With III, in 0.1 M NaCl, the corrosion passes through a low max.; then falls to about the same value as with the unstressed metal. With II, corrosion increases slightly with σ , both in NaCl and in atm. exposure. On the whole, stress has but little effect on the corrosion of these Mg alloys; they are highly resistant to corrosion in exposure to an industrial atm., the loss of tensile strength of 1 mm. thick strips in 2 years having been 5, 16, and 20% only for I, II, and III, resp. The corrosion of stressed Al in 0.3 M HCl increases with σ up to about 20%, then remains const.; in 0.6 M NaCl, it is const. up to about $\sigma = 70\%$, then increases (in 120 days expts.). With Zn in 0.0125 M H₂SO₄, the loss of wt. increases with σ ; the loss of tensile strength first decreases, remains const. between $\sigma = 10$ and 50%, then increases. With Cu in 0.1 M (NH₄)₂SO₄, buffered to pH 2, the corrosion decreases somewhat with increasing σ ; in 0.005 M H₂SO₄, stress has no effect on the corrosion. With steel in 1.6 M H₂SO₄, the corrosion increases with σ only very slightly, more significantly at $\sigma = 70$ -90%. N. T.

9

Corrosional cracking of Electron alloy AZM in various media. H. M. Zatplahji, Dabudy Akad. Nauk S.S.R., 38, 77-8 (1947). The alloy (Mg with 0.5% Al, 1% Zn, 0.5% Mn) was examd. in H_2SO_4 , Na_2CO_3 , and $NaCl-K_2CrO_4$ solns. Small concns. of H_2SO_4 accelerate cracking type of corrosion, possibly by accentuated potential differences between the anodic zones and the rest of the material; in 0.4 N and higher acid general corrosion predominates. In carbonate solns. min. of anodic zones always exceeds that of general corrosion and cracking always takes place. Dichromate accelerates cracking-corrosion by several hundred fold when added to $NaCl$ soln.; the max. effect occurs at about 0.025 M.

Corrosive cracking of magnesium alloys. I. E. M. Zaretskii. J. Applied Chem. (U.S.S.R.) 20, 123-9 (1947) [in Russian]. — Stress-corrosion tests were run on the Mg alloys Electron MA-3 (Al 5.84, Zn 1.10, Mn 0.38, Si 0.14, Fe 0.02), Electron MA-1 (0.15, 0, 1.22, 0.08, 0.01), and Mg (0.33, 0.07, 0.03, 0.02, 0.05), annealed at 420° and etched, in the form of rolled ribbons 1 mm. thick, 15 mm. wide, 100 mm. long, bent into loops of radius 8 mm. and subjected to different known stresses by pulling together or apart the ends of the loop. In atm. exposure, the 1st crack appeared, in MA-3, after 1-2 days under 15.0 kg./sq. mm., after 5-10 days under 0.8, none after 300 days without stress. Similar effects, but after longer times, were observed in immersion in distd. H₂O, and, even more slowly, in a tap-water spray. Under the same conditions, general (non-stress) corrosion was insignificant; nor did any cracking occur in samples bent (plastically) into a loop but not subjected to an elastic stress. In distd. H₂O, in the absence of O₂, the tendency to stress corrosion is decreased markedly. The aspect of stress corrosion is very different from ordinary corrosion; the surface of cracked samples often remained bright. On micrographs, stress-corrosion cracks are seen to pass both along boundaries of grain and through the grains. Annealing followed by cooling under different conditions, and tempering, had little influence on the time of appearance of stress-corrosion cracks. Chem. surface treatments retarded somewhat the appearance of the 1st crack without, however, preventing it seriously. Relatively best results were obtained by anodic oxidation in a soln. K₂Cr₂O₇ 100 + Na₂SiO₃ 50 g./l. Surface treatment with steel balls was of no avail. The MA-1 alloy and Mg are not subject to

stress corrosion. II. Corrosive cracking of cast magnesium alloys. Ibid. 23-6.—(1) In samples 160 X 15 X 3 mm., cut from metal cast in sand at 730-80° and etched, bent under known loads, Mg showed no stress corrosion. Likewise, and in contrast to its behavior in the rolled state, cast MA-3 proved to be resistant; the 1st crack, under 11 kg./sq. mm., appeared in atm. exposure only after 20 days, none in 300 days under smaller stresses. In tap-water spray, the 1st crack appeared, under 11 kg./sq. mm., after 10 months, and only microcracks were found under 0 kg./sq. mm. A much more marked tendency to stress corrosion was found with cast MA-6 (Al 10.0, Zn 0, Mn 0.20, Fe 0.04, Si 0.09); the 1st crack, in atm. exposure, appearing within 2-20 days under 2.5-8.0 kg./sq. mm.; the cracks are mainly along grain boundaries. In tap-water spray, the alloy is highly resistant, no surface cracks being visible in 10 months, only microscopic intercrys. corrosion with crumbling of single crystal grains. Immersion in distd. H₂O causes much more rapid stress corrosion than water spray. Coating with gun grease prevented stress corrosion for 3 years. (2) In further contrast to the rolled alloys, heat-treatment does have a marked effect on the stress corrosion of cast MA-6. Three hrs. annealing at 420°, if followed by cooling in the furnace, lowers the resistance (the 1st crack appearing, in distd. H₂O, in 54 hrs. under 8.2 kg./sq. mm.), but, if followed by quenching in water at 20° or at 110°, it increases the resistance (to 585 hrs.). In samples quenched at 20°, cracking was more pronouncedly intercrys. with crumbling of grains, whereas in samples quenched at 100° the cracks were more nearly oriented in the direction of the stress. The particular sensitivity of the stress-

corrosion resistance of the MA-6 alloy to heat-treatment is linked with the presence of the solid solns. Mg-Mg₂Al₃ and Al-Mg₂Al₃. (ii) An attempt is made to classify stress-corrosion effects in 3 groups. The 1st group comprises alloys with a tendency to intercrys. corrosion even in the absence of deformation; imposition of stress only pulls apart grains already eroded along their boundaries and facilitates further penetration. In the 2nd group, the tendency to corrosion appears as a result of segm. of components from a supersatd. solid soln., along grain boundaries, under the action of the stress (e.g., Al + 6 Mg + 4 Zn, quenched); in this case, homogenization of the grains through annealing should counteract stress corrosion. The 3rd group comprises alloys close to equil. and not subject to intercrys. corrosion in the absence of stress; here, stress corrosion must be attributed mainly to the appearance of local potential differences under the action of the stress.

N. Thon

AGRANAT, B.A.; ZARETSKIY, F.I.

Use of ultrasonic waves to prepare a stable drawing emulsion.
Sbor. nauch. trud. GINTSVETMET no.33:369-376 '60. (MIRA 15:3)
(Metalworking lubricants)
(Ultrasonic waves—Industrial applications)

S/137/62/000/c04/068/201
A052/A101

AUTHORS: Agranat, B. A., Zaretskiy, F. I.

TITLE: Application of ultrasound to the production of a stable drawing emulsion

PERIODICAL: Referativnyy zhurnal, Metallurgiya, no. 4, 1962, 34, abstract 4D197
("Sb. nauchn. tr. In-t tsvet. met. im. M. I. Kalinina", no. 33,
1960. 369-376)

TEXT: The investigation of the emulsification process acted upon by a strong ultrasonic field was carried out on emulsion of the "Moskabel" plant. For drawing Cu- and Al-wire the plant used to prepare emulsions by boiling a soap solution and mechanical mixing with oil and water. These emulsions would usually stratify in 5 - 10 days and proved unsuitable for further use. A series of experiments carried out to produce a more stable emulsion by means of ultrasound has shown that the dispersity of the ultrasound treated emulsion changes inconsiderably, and it preserves practically its stability during several months. A description and diagram of the ultrasonic unit for producing emulsion are given.

K. Ursova

[Abstracter's note: Complete translation]

Card 1/1

KOVALENKO, D.N.; ZARETSKIY, G.M.

Bentonites in the Lopushnianskoye deposit. Bent. gling Ukr.
no.3:3-13 '59.
(MIRA 12:12)

1. Institut geologicheskikh nauk AN USSR i Ukrainskoye
otdeleniye Vsesoyuznogo tresta stroitel'no-tehnicheskikh
izyskaniy.
(Dniester Valley--Bentonite)

SEURKOVICH, S.V.; ZARETSKIY, I.I.

Effect of immunotherapy on kidney function in burns. Khirurgia 35
no.7:16-21 Jl '59. (MIRA 12:12)

1. Iz patofiziologicheskoy laboratorii (zav. - chlen-korrespondent
AMN SSSR prof. N.A. Fedorov) TSentral'nogo instituta geratologii i
perelivaniya krovi (dir. - deystvitel'nyy chlen AMN SSSR prof. A.A.
Bagdasarov) Ministerstva zdравookhraneniya SSSR.

(BURNS, experimental)
(KIDNEYS, physiology)
(IMMUNE SERUMS, pharmacology)

CA ZARETSKII, I. I.

Water metabolism after homogenous blood transfusion.
I. I. Zaretskii (Ministry Health, Moscow). *Arkh. Patol.*
1958, 6, 70-9(1958). Expts. on dogs with fresh citrated
blood (10-18 ml. per kg.) transfusions showed that at first
a drop of H₂O content in the recipient's blood occurs (av.
0.20%), which is followed by a rise, reaching 1.40% (av.)
above normal in 24-48 hrs. Cl level follows that of H₂O.
Erythrocytes participate in the changes involved, as shown
by enhanced ability for uptake of H₂O and Cl within 1-3
days after transfusion; individual vols. of erythrocytes
increase as H₂O accumulation proceeds. Secretory and
secretory processes also go through a decrease-increase
cycle (diuretic and gastric secretion measurements).
Liver tissue acts as a regulator of the changes which take
place. G. M. Kosolapoff

ZARETSKIY, I. I.

35481. Izmeneniye Khimicheskogo i morfologicheskogo sostava Krovi v osle
perelivaniya izogennoy kvovi. Vracheb. delo, 1949, No. 11, stb, 985-88.

Letopis' Zhurnal'nykh Statey, Vol. 48, Moskva, 1949

САРЕПКАЙ, И. Е.
СА

44 G

Water metabolism in healthy and anemic subjects after blood transfusion. I. I. Zaretskii (Ministry of Health, Moscow). *Transfuz. Akad.* 22, No. 3, 65-8(1980); cf. C.I. 44, 5432Y. -The erythrocytes of the recipient take active part in H_2O metabolism in the 1st days after transfusion. They energetically retain H_2O and increase in vol. On the day of transfusion diuresis and elimination of Cl decline, after 24-48 hrs, both rise. Transfusion leads to a temporary (2-3 days) decline in concn. of blood protein (largely globulin). In anemic patients permeability of the capillary endothelium increases. The McClure-Aldrich test is accelerated in the 1st 2-3 days after transfusion.

G. M. Kosolapoff

ZARETSKIY, I. [b] Candidate of Medical Science

"Problems in Connection with Blood Transfusion"

Meditoinskiy Rabotnik, No. 23, 1950

-51-11691, 5 Jul 1950

ZARETSKIY, I. I.

Present problems in hematology and blood transfusion. Sovet.
med. no.9:38-40 Sept 1951. (CIME 21:1)

1. Report on the 29th Expanded Plenary Session of the Learned
Council of the Central Order of Lenin Institute of Hematology
and Blood Transfusion held in May 1951.

ZARETSKIY, I.I., kandidat meditsinskikh nauk (Moscow).

Certain results and prospects in the development of hematology and blood transfusion. Sov.med. 17 no.8:44-47 Ag '53. (MLRA 6:8)
(Blood--Transfusion)

1953 Session of the Scientific Council, Central Order of Lenin
Institute of Hematology and Blood Transfusion

ZARETSKIY, I.I. (Moscow); MIKHAYLOVA, I.A. (Moscow); ROZANOVA, N.S. (Moscow)

Functional state of the kidneys following transfusion of compatible blood. Arkh.pat. 16 no.2:26-31 Ap-Je '54. (MIRA 7:5)

1. Iz patofiziologicheskoy laboratorii (zav. prof. N.A.Fedorov) TSentral'nogo instituta hematologii i perelivaniya krovi, Ministerstva zdravookhraneniya SSSR (dir. chlen-korrespondent AMN SSSR prof. A.A.Bagdasarov).

(BLOOD TRANSFUSION, (KIDNEYS, physiology,
*eff. on kidney funct.) *eff. of blood transfusion)

"APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1

ZARETSKIY, I. I. and FAYNSHTEIN, F. E.

Some New Experimental and Clinical Data on the Field of Hematology.
Voyenno-Meditsinskiy Zhurnal, No 1, p 15, 1955.

APPROVED FOR RELEASE: 09/19/2001

CIA-RDP86-00513R001963820010-1"

FEDOROV, N.A., professor; ZARETSKIY, I.I. (Moskva)

Renal function following blood transfusion; experimental and
clinical investigations. Klin.med., 33 no.11:28-36 N '55.
(MIRA 9:7)

1. Iz patofiziologicheskoy laboratorii (sav.-prof. N.A.Fedorov)
Tsentral'nogo instituta hematologii i perelivaniya krovi
Ministerstva zdravookhraneniya SSSR (dir.-chlen-korrespondent
AMN SSSR prof. A.A.Bagdasarov)
(KIDNEYS, physiology,
eff. of blood transfusion)
(BLOOD TRANSFUSION,
eff. on kidney funct.)

ZARETSKIY, I. I., kandidat meditsinskikh nauk

New trends in the field of hematology and blood transfusion.
Voen.-med. zhur. no.6:8-16 Je '56. (MLRA 9:9)
(BLOOD)

, USSR / Human and Animal Physiology. Excretion.

T

Abs Jour: Ref Zhur-Biol., No 22, 1958, 101947.

Abstract: activity of the tubules according to diodrast or phenol red were determined. The investigations were conducted on the day of transfusion after the emergence of the animals from the state of shock, next day, and later every two days until return to normal. In heterotransfusion and in isotransfusion, a biphasic change of the kidney function occurred, sharper in the first case. During the first phase (1-2 days) after blood infusion, the excretory and concentrating powers of the kidneys decreased, glomerulus filtration and the kidney blood circulation decreased, the filtration fraction and permeability of the glomerular membrane increased. This was confirmed by histologic data (edema of the medullar layer of the kidneys, edema of the glomerular capsule, constriction of the

Card 2/3

ZARETSKIY, I. I.
VINOKUROVA, G.P.; ZARETSKIY, I.I.; MIKHAYLOVA, I.A.

The effect of blood transfusion, blood components and plasma substitutes on kidney functions. Probl.gemat. i perel.krovi 1 no.2: 48-52 Mr-Ap '56. (MIRA 10:1)

1. Iz TSentral'nogo ordena Lenina instituta hematologii i perelivaniya krovi (dir. - chlen-korrespondent AMN SSSR prof. A.A.Badasarov) Ministerstva zdravookhraneniya SSSR.

(KIDNEYS, physiol.

funct., eff. of blood transfusion, blood components and plasma substitutes)

(BLOOD TRANSFUSION

eff. on kidney funct.)

(PLASMA SUBSTITUTES, eff.

on kidney funct.)